

PROPORTION OF ATKIN-LEHNER SIGN PATTERNS AND HECKE EIGENVALUE EQUIDISTRIBUTION

ERICK ROSS, ALEXANDRE VAN LIDTH, MARTHA ROSE WOLF, AND HUI XUE

ABSTRACT. Let $N \geq 1$, $k \geq 2$ even, and σ denote a sign pattern for N . In this paper, we first determine the exact proportion of forms in $S_k(N)$ and $S_k^{\text{new}}(N)$ with a given Atkin-Lehner sign pattern σ . Then we study the asymptotic behavior of the Hecke operators T_p over the subspaces of $S_k(N)$ and $S_k^{\text{new}}(N)$ with Atkin-Lehner sign pattern σ . In particular, for the p -adic Plancherel measure μ_p , we show that the Hecke eigenvalues for T_p over these subspaces are μ_p -equidistributed as $N + k \rightarrow \infty$.

1. INTRODUCTION

For $N \geq 1$ and $k \geq 2$ even, let $S_k(N)$ denote the space of cusp forms of weight k and modular group $\Gamma_0(N)$, and let $S_k^{\text{new}}(N) \subseteq S_k(N)$ denote its new subspace. For $p^r \parallel N$, let W_{p^r} denote the Atkin-Lehner involutions on these spaces, and recall that these W_{p^r} can be extended multiplicatively to W_Q for all $Q \parallel N$ (i.e. $Q|N$ where $(Q, N/Q) = 1$).

Now, recall that the W_Q involutions diagonalize $S_k(N)$ and $S_k^{\text{new}}(N)$ with eigenvalues ± 1 . So, we define the Atkin-Lehner sign pattern spaces by specifying these signs for all $p^r \parallel N$. To be precise, define a sign pattern σ for N to be a multiplicative function on the exact divisors of N such that $\sigma(p^r) = \pm 1$ for all $p^r \parallel N$. Then we have the decomposition of $S_k(N)$

$$S_k(N) = \bigoplus_{\sigma} S_k^{\sigma}(N), \quad S_k^{\sigma}(N) := \{f \in S_k(N) : W_{p^r} f = \sigma(p^r) f \text{ for each } p^r \parallel N\}$$

(and similarly for $S_k^{\text{new}}(N)$). Note that these sign pattern spaces $S_k^{\sigma}(N)$ are a finer decomposition of the more well-known Fricke sign spaces $S_k^{\pm}(N)$, which only specify the “global” sign (i.e. the sign of the Fricke involution W_N).

Now, a natural question one might ask is how all the possible sign patterns are distributed over the spaces $S_k(N)$ and $S_k^{\text{new}}(N)$. For both global signs and local sign patterns, we determine the exact proportion of signs / sign patterns appearing.

In the following theorem, $\mathbb{1}_{\text{condition}}$ denotes the indicator function for *condition*, and the notation “ $A(N, k) \sim B(N)$ as $N + k \rightarrow \infty$ ” means that $A(N, k)/B(N) \rightarrow 1$ as $N + k \rightarrow \infty$. We also take the convention that $0 \sim 0$, because as discussed shortly, both sides of (1.4) are identically zero in one particular case (see Theorem 1.2).

2020 *Mathematics Subject Classification*. Primary 11F25; Secondary 11F72 and 11F11.

Key words and phrases. Hecke operator; Hecke eigenvalues; Eichler-Selberg trace formula; Sign pattern; Equidistribution.

Theorem 1.1. *Consider $N \geq 1$ and $k \geq 2$ even. Then as $N + k \rightarrow \infty$, the proportion of cusp forms with global sign $+1$ is given by*

$$\frac{\dim S_k^+(N)}{\dim S_k(N)} \sim \frac{1}{2} + \frac{\mathbb{1}_{N=1}}{2}, \quad (1.1)$$

$$\frac{\dim S_k^{\text{new},+}(N)}{\dim S_k^{\text{new}}(N)} \sim \frac{1}{2} + \frac{\mathbb{1}_{N=\text{cubefree square}}}{2} \prod_{p|N} \frac{-1}{p^2 - p - 1}. \quad (1.2)$$

Also, consider sign patterns σ for N . Then as $N + k \rightarrow \infty$, the proportion of cusp forms with sign pattern σ is given by

$$\frac{\dim S_k^\sigma(N)}{\dim S_k(N)} \sim \frac{1}{2^{\omega(N)}}, \quad (1.3)$$

$$\frac{\dim S_k^{\text{new},\sigma}(N)}{\dim S_k^{\text{new}}(N)} \sim \frac{1}{2^{\omega(N)}} \prod_{p^r || N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right). \quad (1.4)$$

We note that this theorem predicts an equal proportion of both signs (and more generally, all sign patterns) in many cases. Over the full space, both signs (respectively, all sign patterns) will have equal proportion in $S_k(N)$ for all $N \geq 2$, by (1.1) and (1.3). Over the newspace, both global signs (respectively, all sign patterns) will have equal proportion in $S_k^{\text{new}}(N)$ for squarefree $N \geq 2$, by (1.2) and (1.4). This equiproportion property for the case of newspaces of squarefree level was also shown by Iwaniec-Luo-Sarnak [9, Equation (2.73)] (respectively, Martin [11, Corollary 3.4]).

The paper [13] recently investigated small biases in global sign, and claimed that each global sign would appear for 50% of newforms as $N + k \rightarrow \infty$ [13, Section 1.3]. However, as one can see from (1.2), this is not quite accurate; the global sign in $S_k^{\text{new}}(N)$ will actually be asymptotically biased towards $(-1)^{\omega(N)}$ whenever N is a cubefree square. One can see this bias numerically from the data calculated in [18].

We would also like to point out that Martin [14] recently investigated small biases in the sign of W_{p^r} for a single fixed prime power p^r , and observed a stronger bias at $r = 2$. Here, Theorem 1.1 quantifies this observation by establishing the exact proportion of each sign pattern.

Now, observe that the proportions (1.3) and (1.4) are always nonzero, except when $4 || N$ with $\sigma(4) = +1$ over the newspace $S_k^{\text{new}}(N)$. For this reason, we define the notion of “admissible” sign patterns, where all sign patterns are admissible except for the one case of $4 || N$ with $\sigma(4) = +1$ over the newspace. It is worth noting here that the admissible sign patterns σ all have proportion within a factor of 3 of $1/2^{\omega(N)}$. In other words, the admissible sign patterns are not too far off from being equally proportioned. This “factor of 3” comes from the lower and upper bounds on the $\prod_{p^r || N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right)$ factor from (1.4) for admissible σ :

$$\begin{aligned} 0.71546 &\leq \prod_{p \neq 2} \left(1 - \frac{1}{p^2 - p - 1} \right) \leq \prod_{p^r || N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right) \\ &\leq \prod_p \left(1 + \frac{1}{p^2 - p - 1} \right) \leq 2.67412. \end{aligned} \quad (1.5)$$

It turns out that for inadmissible sign patterns σ , there are no newforms at all with sign pattern σ . This is because W_4 always has Atkin-Lehner sign -1 over $S_k^{\text{new}}(N)$. This result was originally shown by Atkin and Lehner in [2, Theorem 7]. We give an alternative proof here based on the trace formula.

Theorem 1.2. *Let $N \geq 1$ with $4 \parallel N$. Then $W_4 f = -f$ for every $f \in S_k^{\text{new}}(N)$.*

Now, recall that the Hecke operators T_m for $(m, N) = 1$ commute with the Atkin-Lehner involutions W_Q [6, Proposition 13.2.6] (see [6] for the precise definition of these Hecke operators T_m). This means that the Hecke operators preserve the sign pattern spaces $S_k^\sigma(N)$ and $S_k^{\text{new}, \sigma}(N)$. So for the admissible sign patterns σ , we can then investigate how the Hecke eigenvalues over $S_k^\sigma(N)$ and $S_k^{\text{new}, \sigma}(N)$ are distributed. For $m \geq 1$, let $T'_m := m^{-(k-1)/2} T_m$ denote the normalized Hecke operators (over both $S_k^\sigma(N)$ and $S_k^{\text{new}, \sigma}(N)$). Then for a fixed prime p , let $\mu_p(x) := \frac{p+1}{\pi} \frac{(1-x^2/4)^{1/2}}{(p^{1/2} + p^{-1/2})^2 - x^2} dx$ denote the p -adic Plancherel measure. We will show that the eigenvalues of T'_p (over both $S_k^\sigma(N)$ and $S_k^{\text{new}, \sigma}(N)$) are μ_p -equidistributed as $N+k \rightarrow \infty$.

Theorem 1.3. *Fix p prime. Consider $N \geq 1$ coprime to p , σ sign patterns for N , and $k \geq 2$ even. Then the Hecke eigenvalues of $T'_p|_{S_k^\sigma(N)}$ are μ_p -equidistributed as $N+k \rightarrow \infty$. In other words, for any sequence $\{(N_i, k_i, \sigma_i)\}_{i \geq 1}$ with $N_i + k_i \rightarrow \infty$, the collection of Hecke eigenvalues $\text{eigv}(T'_m|_{S_{k_i}^{\sigma_i}(N_i)})$ is μ_p -equidistributed as $i \rightarrow \infty$. Moreover, if we assume it is not the case that $4 \parallel N$ with $\sigma(4) = +1$, then the Hecke eigenvalues of $T'_p|_{S_k^{\text{new}, \sigma}(N)}$ are also μ_p -equidistributed as $N+k \rightarrow \infty$.*

Note here that similar μ_p -equidistribution results have been shown before in different settings. For example, Conrey-Duke-Farmer showed μ_p -equidistribution over $S_k(1)$ [7] and Serre showed μ_p -equidistribution over $S_k(N)$, $S_k^{\text{new}}(N)$, and $S_k(N, \chi)$ [19]. More recently, the first author showed μ_p -equidistribution over $S_k^{\text{new}}(N, \chi)$ for admissible characters χ [17]. Theorem 1.3 can be interpreted as strengthening Serre's result over $S_k(N)$ and $S_k^{\text{new}}(N)$, showing that μ_p -equidistribution still holds over the much finer sign pattern decompositions $S_k^\sigma(N)$ and $S_k^{\text{new}, \sigma}(N)$. In fact, this sign pattern decomposition is, in some sense, the finest (natural) decomposition possible; the $S_k^{\text{new}, \sigma}(N)$ seem to be the smallest subspaces of $S_k^{\text{new}}(N)$ that still respect the Hecke operators and Galois conjugation [4, Section 5].

Lastly, in Section 6, we discuss several applications of Theorem 1.3, similar to those in [19, §6]. The μ_p -equidistribution shown in Theorem 1.3 can be used to obtain information about the Galois orbits of Hecke eigenvalues, or equivalently, about the coefficient fields $\mathbb{Q}(f)$ of newforms $f \in S_k^{\text{new}, \sigma}(N)$. In particular, we show that the degrees of these coefficient fields are unbounded as $N \rightarrow \infty$. This can be interpreted in terms of a weak form of the generalized Maeda conjecture. We are also able to obtain information about the simple \mathbb{Q} -factors of the Jacobian of the modular curve $X_0(N)$. See Section 6 for precise details.

We now give an overview of the arguments used in this paper. In Section 2, we use a trace formula from Skoruppa-Zagier [20] to estimate $\text{Tr } T'_m \circ W_Q$ over both $S_k(N)$ and $S_k^{\text{new}}(N)$. Then in

Section 3, we use the results of Section 2 to obtain a trace formula for T'_m over the sign pattern spaces. Next, in Section 4, we prove Theorems 1.1 and 1.2, computing the exact proportion of how often each sign pattern can appear. Then, in Section 5, we use the $\text{Tr } T'_m$ estimates of Section 3 to prove the μ_p -equidistribution of the Hecke eigenvalues over $S_k^\sigma(N)$ and $S_k^{\text{new},\sigma}(N)$. In this section, we use the same general strategy used by Serre [19]. Finally, in Section 6, we discuss applications of Theorem 1.3.

2. TRACE FORMULA FOR $T'_m \circ W_Q$

For integers $k \geq 2$, let $p_k(t, m)$ denote the Lucas sequence of the first kind; i.e. $p_k(t, m)$ is the x^{k-2} coefficient in the power series expansion of $(mx^2 - tx + 1)^{-1}$. Note that in particular, we have by Clayton et al. [5, Lemmas 2.2, 2.3] that

$$\begin{cases} p_k(t, m) = (k-1)m^{(k-2)/2} & \text{if } t^2 - 4m = 0, \\ p_k(t, m) \leq \frac{2m^{(k-1)/2}}{\sqrt{|t^2 - 4m|}} & \text{if } t^2 - 4m < 0. \end{cases} \quad (2.1)$$

Following Skoruppa-Zagier [20], we also define the following class numbers. For integers $N \geq 1$ and $\Delta \leq 0$, define $H_N(\Delta)$ via

$$H_N(\Delta) = \begin{cases} a^2b \left(\frac{\Delta/a^2b^2}{N/a^2b} \right) H(|\Delta/a^2b^2|) & \text{if } a^2b^2 \mid \Delta, \\ 0 & \text{otherwise,} \end{cases} \quad (2.2)$$

where $a^2b := (N, \Delta)$ with b squarefree, (\cdot) denotes the Kronecker symbol, and $H(\cdot)$ denotes the Hurwitz-Kronecker class number. Note that in the case of $\Delta = 0$, $\left(\frac{0}{1}\right) = 1$ and $H(0) = \frac{-1}{12}$, so that $H_N(0) = \frac{-1}{12}N$.

In the following, we use the notation $Q \parallel N$ to mean $Q \mid N$ and $(Q, \frac{N}{Q}) = 1$ and $\ell = \square$ to mean the integer ℓ is a square. In addition, $\sigma_0(N) = \sum_{N' \mid N} 1$, $\sigma_1(N) = \sum_{N' \mid N} N'$, and $B(N)$ is defined to be the greatest integer ℓ such that $\ell^2 \mid N$.

Proposition 2.1 (Skoruppa-Zagier [20], correction by Assaf [1]). *For $N \geq 1$ coprime to m , $Q \parallel N$, and $k \geq 2$ even, we have*

$$\begin{aligned} \text{Tr}_{S_k(N)} T'_m \circ W_Q &= \sum_{\substack{N' \mid N \\ N/N' \text{ squarefree}}} \mu \left(\frac{Q}{(Q, N')} \right) s'_{k, N'}(m, (Q, N')), \\ \text{Tr}_{S_k^{\text{new}}(N)} T'_m \circ W_Q &= \sum_{N' \mid N} \alpha \left(\frac{N}{N'} \right) s'_{k, N'}(m, (Q, N')). \end{aligned}$$

Here μ denotes the Möbius function, $\alpha(N)$ is the nonzero multiplicative function defined on prime powers via

$$\alpha(p^r) = \begin{cases} -1 & r = 1 \text{ or } 2 \\ 1 & r = 3 \\ 0 & r \geq 4, \end{cases}$$

and

$$s'_{k,N}(m, Q) = -\frac{1}{2m^{(k-1)/2}} \sum_{Q'|Q} \sum_s p_k\left(\frac{s}{\sqrt{Q'}}, m\right) H_{\frac{N}{Q}}(s^2 - 4mQ') \quad (2.3)$$

$$- \frac{1}{2m^{(k-1)/2}} \sum_{m'|m} \min\left(m', \frac{m}{m'}\right)^{k-1} \left(B(Q), m' + \frac{m}{m'}\right) \left(B\left(\frac{N}{Q}\right), m' - \frac{m}{m'}\right) \quad (2.4)$$

$$+ \frac{1}{m^{(k-1)/2}} \mathbb{1}_{\substack{k=2 \\ \frac{N}{Q}=\square}} \sigma_0(Q) \sigma_1(m), \quad (2.5)$$

with the summation over s ranging over all $s \in \mathbb{Z}$ such that $s^2 \leq 4mQ'$, $Q'|s$, and $\left(\left(\frac{s}{Q'}\right)^2, \frac{Q}{Q'}\right)$ is squarefree.

We then estimate the $s'_{k,N}(m, Q)$ terms from this proposition. In the following, and throughout the entire paper, we use big- O notation with respect to both N and k . Note that m here is a fixed constant (so the big- O implied constants will depend on m). And Q is considered to be an arbitrary exact divisor of N (so the big- O bounds will only be stated in terms of N and k , uniform over arbitrary $Q \parallel N$).

Lemma 2.2. Fix $m \geq 1$. Let $N \geq 1$ be coprime to m , $Q \parallel N$, and $k \geq 2$ be even. Then

$$s'_{k,N}(m, Q) = \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \frac{N}{Q} + O(N^{1/2} \sigma_0(N) \log N),$$

where $s'_{k,N}(m, Q)$ is as defined in Proposition 2.1.

Proof. Note that $\sigma_0(Q) \leq \sigma_0(N) = O(N^\varepsilon)$ and $\sigma_1(m) = O(1)$, so the third term (2.5) of $s'_{k,N}(m, Q)$ is $O(N^\varepsilon)$. Also note that since

$$\left(B(Q), m' + \frac{m}{m'}\right) \leq 2m, \quad \left(B\left(\frac{N}{Q}\right), m' - \frac{m}{m'}\right) \leq \sqrt{\frac{N}{Q}} \leq \sqrt{N},$$

the second term (2.4) of $s'_{k,N}(m, Q)$ is of magnitude

$$\leq \frac{1}{2m^{(k-1)/2}} \sum_{m'|m} \min\left(m', \frac{m}{m'}\right)^{k-1} (2m) \sqrt{N} \leq \frac{(2m) \sqrt{N}}{2} \sum_{m'|m} 1 = O(\sqrt{N}).$$

Lastly, we consider the first term (2.3) of $s'_{k,N}(m, Q)$. In the double summation, denote $\Delta := s^2 - 4mQ'$, and we break into two cases: when $\Delta < 0$, and when $\Delta = 0$.

When $\Delta < 0$, we have by (2.1) that

$$\left| p_k \left(\frac{s}{\sqrt{Q'}}, m \right) \right| \leq \frac{2m^{(k-1)/2}}{\sqrt{\left| \frac{s^2}{Q'} - 4m \right|}} = \frac{2m^{(k-1)/2}}{\sqrt{|\Delta|/Q'}}.$$

Using [8, Lemma 2.2], we can bound the Hurwitz-Kronecker class number by

$$H(D) \leq \frac{\sqrt{D}(\log D + 2)}{\pi}.$$

So letting $a^2b = \left(\frac{N}{Q}, \Delta \right)$ we have

$$\begin{aligned} \left| H_{\frac{N}{Q}}(\Delta) \right| &\leq \left| a^2b \left(\frac{\Delta/a^2b^2}{N/a^2b} \right) H(|\Delta/a^2b^2|) \right| \\ &\leq \frac{a^2b\sqrt{|\Delta/a^2b^2|}(\log |\frac{\Delta}{a^2b^2}| + 2)}{\pi} \\ &\leq \frac{a\sqrt{|\Delta|}(\log |\Delta| + 2)}{\pi}. \end{aligned}$$

Combining these,

$$\begin{aligned} \left| p_k \left(\frac{s}{\sqrt{Q'}}, m \right) H_{\frac{N}{Q}}(\Delta) \right| &\leq \frac{2m^{(k-1)/2}}{\sqrt{\Delta/Q'}} \frac{a\sqrt{|\Delta|}(\log |\Delta| + 2)}{\pi} \\ &= \frac{2m^{(k-1)/2}a\sqrt{Q'}(\log |\Delta| + 2)}{\pi} \\ &\leq \frac{2m^{(k-1)/2}\sqrt{N}(\log(4mN) + 2)}{\pi} \\ &= m^{(k-1)/2}O(N^{1/2} \log N), \end{aligned}$$

where the last inequality comes from the facts that $a^2 \leq \frac{N}{Q} \leq \frac{N}{Q'}$ and $|\Delta| \leq 4mQ' \leq 4mN$.

Now, note that in (2.3), the number of terms in the summation over $Q'|Q$ is equal to $\sigma_0(Q) \leq \sigma_0(N)$. Furthermore, in (2.3), the number of terms in the summation over s is $\leq 4\sqrt{m/Q'} = O(1)$ since $Q'|s$ and $|s| \leq 2\sqrt{mQ'}$. Thus, the contribution of the $\Delta < 0$ terms in $s'_{k,N}(m, Q)$ is $O(N^{1/2}\sigma_0(N) \log N)$.

Finally, we consider the $\Delta = 0$ terms, which will make up the main term of $s'_{k,N}(m, Q)$. Recall that $(m, Q') = 1$. Then note that $\Delta := s^2 - 4mQ' = 0$ implies that $m = \square$ and $Q' = \square$. Additionally, $(Q')^2 | s^2 = 4mQ'$ implies that $Q' | 4$. Hence we must have $Q' = 1$ or 4 . It is straightforward to verify that the first case occurs precisely when $m = \square$ and $4 \nmid Q$ (for $Q' = 1$, $s = \pm 2\sqrt{m}$); and the second case occurs precisely when $m = \square$ and $4|Q$ (for $Q' = 4$, $s = \pm 4\sqrt{m}$). This means that in all cases, the number of $\Delta = 0$ terms appearing in $s'_{k,N}(m, Q)$ is exactly $2 \mathbb{1}_{m=\square}$.

And when $\Delta = 0$, we have by (2.1) and (2.2) that

$$p_k \left(\frac{s}{\sqrt{Q'}}, m \right) = (k-1) m^{(k-2)/2}$$

$$\text{and } H_{\frac{N}{Q}}(s^2 - 4mQ') = H_{\frac{N}{Q}}(0) = \frac{-1}{12} \frac{N}{Q}.$$

Thus, extracting the $\Delta = 0$ terms from $s'_{k,N}(m, Q)$, we obtain

$$\begin{aligned} & -\frac{1}{2m^{(k-1)/2}} \sum_{Q'|Q} \sum_{\substack{s^2=4mQ' \\ Q'|s}} p_k\left(\frac{s}{\sqrt{Q'}}, m\right) H_{\frac{N}{Q}}(s^2 - 4mQ') \\ &= -\frac{1}{2m^{(k-1)/2}} \left(2\mathbb{1}_{m=\square}\right) \left((k-1)m^{(k-2)/2}\right) \left(\frac{-1}{12} \frac{N}{Q}\right) \\ &= \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \frac{N}{Q}, \end{aligned}$$

completing the proof. \square

With this estimate of $s'_{k,N}(m, Q)$, we can then estimate $\text{Tr } T'_m \circ W_Q$. In the following, we use the notation $\llbracket A \rrbracket_{\text{condition}}$ to denote that the term A only appears when *condition* is satisfied.

Proposition 2.3. *Fix $m \geq 1$. Let $N \geq 1$ be coprime to m , $Q \parallel N$, and $k \geq 2$ be even. Then*

$$\begin{aligned} \text{Tr}_{S_k(N)} T'_m \circ W_Q &= \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \psi(N) \mathbb{1}_{Q=1} + O(N^{1/2} \sigma_0(N)^2 \log N), \\ \text{Tr}_{S_k^{\text{new}}(N)} T'_m \circ W_Q &= \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \psi^{\text{new}}(N) \eta(Q) + O(N^{1/2} \sigma_0(N)^2 \log N), \end{aligned}$$

where ψ , ψ^{new} , and η are the nonzero multiplicative functions defined on prime powers via

$$\begin{aligned} \psi(p^r) &:= p^r \left(1 + \frac{1}{p}\right), \\ \psi^{\text{new}}(p^r) &:= p^r \left(1 - \frac{1}{p} - \left\llbracket \frac{1}{p^2} \right\rrbracket_{r \geq 2} + \left\llbracket \frac{1}{p^3} \right\rrbracket_{r \geq 3}\right), \\ \eta(p^r) &:= \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1}. \end{aligned}$$

Note in particular that when N is squarefree, $\eta(Q) = \mathbb{1}_{Q=1}$.

Proof. Define the multiplicative function $\theta_M(d) := \frac{d}{(M, d)}$. Observe that $\mu \circ \theta_M$ is also multiplicative since $\theta_M(d_1), \theta_M(d_2)$ are coprime whenever d_1, d_2 are coprime.

To show the first desired identity, it suffices (by Lemma 2.2 and Proposition 2.1) to show that

$$\sum_{\substack{N'|N \\ N/N' \text{ squarefree}}} \mu\left(\frac{Q}{(Q, N')}\right) \frac{N'}{(Q, N')} = \psi(N) \mathbb{1}_{Q=1}. \quad (2.6)$$

Observe that $\mathbb{1}_{N/N' \text{ squarefree}} = \mu^2(N/N')$, $\mu\left(\frac{Q}{(Q, N')}\right) = \mu\left(\frac{N/N'}{(N/N', N/Q)}\right) = \mu \circ \theta_{N/Q}(N/N')$, and $\frac{N'}{(Q, N')} = \theta_Q(N')$. This means that

$$\text{LHS of (2.6)} = [[\mu^2 \cdot \mu \circ \theta_{N/Q}] * \theta_Q](N)$$

$$\begin{aligned}
&= \prod_{p^r \parallel N} [[\mu^2 \cdot \mu \circ \theta_{N/Q}] * \theta_Q](p^r) \\
&= \prod_{p^r \parallel N} \sum_{i=0}^r \mu^2(p^i) \mu \circ \theta_{N/Q}(p^i) \theta_Q(p^{r-i}) \\
&= \prod_{p^r \parallel N} \left(\theta_Q(p^r) + \mu \circ \theta_{N/Q}(p) \theta_Q(p^{r-1}) \right) \\
&= \prod_{p^r \parallel N} \begin{cases} p^r + p^{r-1} & \text{if } p \nmid Q \\ 0 & \text{if } p \mid Q \end{cases} \\
&= \psi(N) \mathbb{1}_{Q=1}.
\end{aligned}$$

Note that here, the second-to-last equality comes from the facts that

$$\theta_Q(p^s) = \begin{cases} p^s & \text{if } p \nmid Q \\ 1 & \text{if } p \mid Q \end{cases} \quad \text{and} \quad \mu \circ \theta_{N/Q}(p) = \begin{cases} 1 & \text{if } p \nmid Q \\ -1 & \text{if } p \mid Q. \end{cases}$$

Similarly, to show the second desired identity, it suffices (by Lemma 2.2 and Proposition 2.1) to show that

$$\sum_{N' \mid N} \alpha \left(\frac{N}{N'} \right) \frac{N'}{(Q, N')} = \psi^{\text{new}}(N) \eta(Q). \quad (2.7)$$

And here,

$$\begin{aligned}
\text{LHS of (2.7)} &= [\alpha * \theta_Q](N) \\
&= \prod_{p^r \parallel N} [\alpha * \theta_Q](p^r) \\
&= \prod_{p^r \parallel N} \sum_{i=0}^r \alpha(p^i) \theta_Q(p^{r-i}) \\
&= \prod_{p^r \parallel N} \left(\theta_Q(p^r) - \theta_Q(p^{r-1}) - \llbracket \theta_Q(p^{r-2}) \rrbracket_{r \geq 2} + \llbracket \theta_Q(p^{r-3}) \rrbracket_{r \geq 3} \right) \\
&= \prod_{p^r \parallel N} \begin{cases} \psi^{\text{new}}(p^r) & \text{if } p \nmid Q \\ -\mathbb{1}_{r=2} & \text{if } p \mid Q \end{cases} \\
&= \psi^{\text{new}}(N) \prod_{p \mid Q} \frac{-\mathbb{1}_{v_p(N)=2}}{\psi^{\text{new}}(p^2)} \\
&= \psi^{\text{new}}(N) \prod_{p^r \parallel Q} \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \\
&= \psi^{\text{new}}(N) \eta(Q),
\end{aligned}$$

as desired. □

3. TRACE FORMULA FOR T'_m OVER THE SIGN PATTERN SPACES

In [11, Proposition 3.2], Martin derived a formula for $\dim S_k^{\text{new},\sigma}(N)$ in terms of the traces $\text{Tr}_{S_k^{\text{new}}(N)} W_Q$. Here, we prove a similar formula for $\text{Tr}_{S_k^\sigma(N)} T'_m$ and $\text{Tr}_{S_k^{\text{new},\sigma}(N)} T'_m$ in terms of the traces $\text{Tr}_{S_k(N)} T'_m \circ W_Q$ and $\text{Tr}_{S_k^{\text{new}}(N)} T'_m \circ W_Q$, respectively.

We note that the identity (3.1) underlying this lemma can also be interpreted as a special case of orthogonality for characters over a finite group. (The set of all exact divisors of N forms a group via the binary operation $Q_1 * Q_2 := \frac{Q_1 Q_2}{(Q_1, Q_2)^2}$. And sign patterns for N make up the characters on this group.)

Lemma 3.1. *For $N \geq 1$,*

$$\begin{aligned}\text{Tr}_{S_k^\sigma(N)} T'_m &= \frac{1}{2^{\omega(N)}} \sum_{Q||N} \sigma(Q) \text{Tr}_{S_k(N)} T'_m \circ W_Q, \\ \text{Tr}_{S_k^{\text{new},\sigma}(N)} T'_m &= \frac{1}{2^{\omega(N)}} \sum_{Q||N} \sigma(Q) \text{Tr}_{S_k^{\text{new}}(N)} T'_m \circ W_Q.\end{aligned}$$

Proof. We show this result just for the full space $S_k(N)$, and the argument for the newspace $S_k^{\text{new}}(N)$ is identical.

First, observe that

$$\begin{aligned}\sum_{Q||N} \sigma(Q) \sigma'(Q) &= \prod_{p^r || N} \sum_{Q||p^r} \sigma(Q) \sigma'(Q) = \prod_{p^r || N} (1 + \sigma(p^r) \sigma'(p^r)) \\ &= \prod_{p^r || N} 2 \mathbb{1}_{\sigma(p^r) = \sigma'(p^r)} = 2^{\omega(N)} \mathbb{1}_{\sigma = \sigma'}.\end{aligned}\tag{3.1}$$

Then, proceeding by splitting $\text{Tr}_{S_k(N)} T'_m \circ W_Q$ into a sum of $\text{Tr}_{S_k^{\sigma'}(N)} T'_m \circ W_Q$ for all sign patterns σ' of N :

$$\begin{aligned}\frac{1}{2^{\omega(N)}} \sum_{Q||N} \sigma(Q) \text{Tr}_{S_k(N)} T'_m \circ W_Q &= \frac{1}{2^{\omega(N)}} \sum_{Q||N} \sigma(Q) \sum_{\sigma'} \text{Tr}_{S_k^{\sigma'}(N)} T'_m \circ W_Q \\ &= \frac{1}{2^{\omega(N)}} \sum_{Q||N} \sum_{\sigma'} \sigma(Q) \sigma'(Q) \text{Tr}_{S_k^{\sigma'}(N)} T'_m \\ &= \frac{1}{2^{\omega(N)}} \sum_{\sigma'} 2^{\omega(N)} \mathbb{1}_{\sigma = \sigma'} \text{Tr}_{S_k^{\sigma'}(N)} T'_m \\ &= \text{Tr}_{S_k^\sigma(N)} T'_m,\end{aligned}$$

as desired. \square

We can then estimate $\text{Tr}_{S_k^\sigma(N)} T'_m$ and $\text{Tr}_{S_k^{\text{new},\sigma}(N)} T'_m$ by applying Proposition 2.3 to Lemma 3.1.

Corollary 3.2. *Fix $m \geq 1$. Then for $N \geq 1$ coprime to m and $k \geq 2$ even,*

$$\text{Tr}_{S_k^\sigma(N)} T'_m = \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \frac{\psi(N)}{2^{\omega(N)}} + O(N^{1/2} \sigma_0(N)^2 \log N),\tag{3.2}$$

$$\mathrm{Tr}_{S_k^{\mathrm{new}, \sigma}(N)} T'_m = \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \frac{\psi^{\mathrm{new}}(N)}{2^{\omega(N)}} \prod_{p^r \parallel N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right) + O(N^{1/2} \sigma_0(N)^2 \log N). \quad (3.3)$$

Proof. Using Lemma 3.1 and then Proposition 2.3, we obtain

$$\begin{aligned} \mathrm{Tr}_{S_k^\sigma(N)} T'_m &= \frac{1}{2^{\omega(N)}} \sum_{Q \parallel N} \sigma(Q) \mathrm{Tr}_{S_k(N)} T'_m \circ W_Q \\ &= \frac{1}{2^{\omega(N)}} \sum_{Q \parallel N} \sigma(Q) \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \psi(N) \mathbb{1}_{Q=1} + O(N^{1/2} \sigma_0(N)^2 \log N) \\ &= \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \frac{\psi(N)}{2^{\omega(N)}} + O(N^{1/2} \sigma_0(N)^2 \log N), \\ \mathrm{Tr}_{S_k^{\mathrm{new}, \sigma}(N)} T'_m &= \sum_{Q \parallel N} \sigma(Q) \mathrm{Tr}_{S_k^{\mathrm{new}}(N)} T'_m \circ W_Q \\ &= \frac{1}{2^{\omega(N)}} \sum_{Q \parallel N} \sigma(Q) \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \psi^{\mathrm{new}}(N) \eta(Q) + O(N^{1/2} \sigma_0(N)^2 \log N) \\ &= \frac{\mathbb{1}_{m=\square}}{\sqrt{m}} \frac{k-1}{12} \frac{\psi^{\mathrm{new}}(N)}{2^{\omega(N)}} \prod_{p^r \parallel N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right) + O(N^{1/2} \sigma_0(N)^2 \log N), \end{aligned}$$

as desired. \square

We point out here that the main term of (3.2) always grows faster than the error term $O(N^{1/2} \sigma_0(N)^2 \log N) = O(N^{1/2+\varepsilon})$. Furthermore, the main term of (3.3) grows faster than the error term, assuming it is not the case that $4 \parallel N$ with $\sigma(4) = +1$ (because after excluding this one particular case, the $\prod_{p^r \parallel N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right)$ factor in (3.3) is $\geq 1/2$ by (1.5)). In other words, the main terms of this corollary grow faster than the corresponding error terms for all admissible sign patterns σ .

4. PROPORTION OF SIGN PATTERNS

In this section, we show Theorems 1.1 and 1.2.

Theorem 1.1. *Consider $N \geq 1$ and $k \geq 2$ even. Then as $N + k \rightarrow \infty$, the proportion of cusp forms with global sign $+1$ is given by*

$$\begin{aligned} \frac{\dim S_k^+(N)}{\dim S_k(N)} &\sim \frac{1}{2} + \frac{\mathbb{1}_{N=1}}{2}, \\ \frac{\dim S_k^{\mathrm{new}, +}(N)}{\dim S_k^{\mathrm{new}}(N)} &\sim \frac{1}{2} + \frac{\mathbb{1}_{N=\text{cubefree square}}}{2} \prod_{p \mid N} \frac{-1}{p^2 - p - 1}. \end{aligned}$$

Moreover, consider sign patterns σ for N . Then as $N + k \rightarrow \infty$, the proportion of cusp forms with sign pattern σ is given by

$$\frac{\dim S_k^\sigma(N)}{\dim S_k(N)} \sim \frac{1}{2^{\omega(N)}}, \quad (4.1)$$

$$\frac{\dim S_k^{\text{new},\sigma}(N)}{\dim S_k^{\text{new}}(N)} \sim \frac{1}{2^{\omega(N)}} \prod_{p^r \parallel N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right). \quad (4.2)$$

Proof. First, substitute $m = 1$ and $Q = 1$ into Proposition 2.3 to obtain the dimensions of the entire spaces:

$$\dim S_k(N) = \frac{k-1}{12} \psi(N) + O(N^{1/2+\varepsilon}), \quad (4.3)$$

$$\dim S_k^{\text{new}}(N) = \frac{k-1}{12} \psi^{\text{new}}(N) + O(N^{1/2+\varepsilon}). \quad (4.4)$$

Now, for the first two claims, we use Proposition 2.3 at $m = 1$ and $Q = 1, N$ to obtain

$$\begin{aligned} \dim S_k^+(N) &= \frac{1}{2} (\text{Tr}_{S_k(N)} W_1 + \text{Tr}_{S_k(N)} W_N) \\ &= \frac{k-1}{12} \psi(N) \frac{1 + \mathbb{1}_{N=1}}{2} + O(N^{1/2+\varepsilon}), \\ \dim S_k^{\text{new},+}(N) &= \frac{1}{2} (\text{Tr}_{S_k^{\text{new}}(N)} W_1 + \text{Tr}_{S_k^{\text{new}}(N)} W_N) \\ &= \frac{k-1}{12} \psi^{\text{new}}(N) \frac{1 + \eta(N)}{2} + O(N^{1/2+\varepsilon}) \\ &= \frac{k-1}{12} \psi^{\text{new}}(N) \frac{1 + \mathbb{1}_{N=\text{cubefree square}} \prod_{p|N} \frac{-1}{p^2 - p - 1}}{2} + O(N^{1/2+\varepsilon}). \end{aligned}$$

Comparing these estimates with (4.3) and (4.4) then yields the desired result.

For the last two claims, we substitute $m = 1$ into Corollary 3.2 to obtain

$$\begin{aligned} \dim S_k^\sigma(N) &= \frac{1}{2^{\omega(N)}} \frac{k-1}{12} \psi(N) + O(N^{1/2+\varepsilon}), \\ \dim S_k^{\text{new},\sigma}(N) &= \frac{1}{2^{\omega(N)}} \frac{k-1}{12} \psi^{\text{new}}(N) \prod_{p^r \parallel N} \left(1 + \sigma(p^r) \frac{-\mathbb{1}_{r=2}}{p^2 - p - 1} \right) + O(N^{1/2+\varepsilon}). \end{aligned}$$

Comparing these estimates with (4.3) and (4.4) similarly yields the desired result. \square

Recall that we defined the notion of admissible sign patterns, where all sign patterns are admissible except for the case of $4 \parallel N$ with $\sigma(4) = +1$ over the newspace. We introduced this definition because the proportions (4.1) and (4.2) are always nonzero except in this one exceptional case.

In fact, it turns out that W_4 always has Atkin-Lehner sign -1 over $S_k^{\text{new}}(N)$. This means that for inadmissible sign patterns σ , there are no forms at all in $S_k^{\text{new}}(N)$ with sign pattern σ .

Theorem 1.2. *Let $N \geq 1$ with $4 \parallel N$. Then $W_4 f = -f$ for every $f \in S_k^{\text{new}}(N)$.*

Proof. It suffices to show that $\text{Tr}_{S_k^{\text{new}}(N)} W_1 + \text{Tr}_{S_k^{\text{new}}(N)} W_4 = 0$. Using Proposition 2.1, we compute

$$\begin{aligned} &\text{Tr}_{S_k^{\text{new}}(N)} W_1 + \text{Tr}_{S_k^{\text{new}}(N)} W_4 \\ &= \sum_{N' \mid N} \alpha \left(\frac{N}{N'} \right) \left[s'_{k,N'}(1, (1, N')) + s'_{k,N'}(1, (4, N')) \right] \end{aligned}$$

$$\begin{aligned}
&= \sum_{L|\frac{N}{4}} \sum_{N' \in \{L, 2L, 4L\}} \alpha\left(\frac{N}{N'}\right) \left[s'_{k,N'}(1, (1, N')) + s'_{k,N'}(1, (4, N')) \right] \\
&= \sum_{L|\frac{N}{4}} \left[\alpha\left(\frac{N}{L}\right) \left[s'_{k,L}(1, 1) + s'_{k,L}(1, 1) \right] + \alpha\left(\frac{N}{2L}\right) \left[s'_{k,2L}(1, 1) + s'_{k,2L}(1, 2) \right] \right. \\
&\quad \left. + \alpha\left(\frac{N}{4L}\right) \left[s'_{k,4L}(1, 1) + s'_{k,4L}(1, 4) \right] \right] \\
&= \sum_{L|\frac{N}{4}} \alpha\left(\frac{N}{4L}\right) \left[-2s'_{k,L}(1, 1) - s'_{k,2L}(1, 1) - s'_{k,2L}(1, 2) + s'_{k,4L}(1, 1) + s'_{k,4L}(1, 4) \right] \\
&=: \sum_{L|\frac{N}{4}} \alpha\left(\frac{N}{4L}\right) [*].
\end{aligned}$$

Now, we show that $[*] = 0$. Computing each of the terms appearing in $[*]$ (see (2.3)-(2.5) for the definition of $s'_{k,L}$) yields

$$\begin{aligned}
s'_{k,L}(1, 1) &= -\frac{1}{2} \left[p_k(0, 1)H_L(-4) + 2p_k(1, 1)H_L(-3) + 2p_k(2, 1)H_L(0) + B(L) - 2\mathbb{1}_{\substack{k=2 \\ L=\square}} \right], \\
s'_{k,2L}(1, 1) &= -\frac{1}{2} \left[p_k(0, 1)H_{2L}(-4) + 2p_k(1, 1)H_{2L}(-3) + 2p_k(2, 1)H_{2L}(0) + B(L) \right], \\
s'_{k,2L}(1, 2) &= -\frac{1}{2} \left[p_k(0, 1)H_L(-4) + 2p_k(1, 1)H_L(-3) + 2p_k(2, 1)H_L(0) + p_k(0, 1)H_L(-8) \right. \\
&\quad \left. + 2p_k(\sqrt{2}, 1)H_L(-4) + B(L) - 4\mathbb{1}_{\substack{k=2 \\ L=\square}} \right], \\
s'_{k,4L}(1, 1) &= -\frac{1}{2} \left[p_k(0, 1)H_{4L}(-4) + 2p_k(1, 1)H_{4L}(-3) + 2p_k(2, 1)H_{4L}(0) + 2B(L) - 2\mathbb{1}_{\substack{k=2 \\ L=\square}} \right], \\
s'_{k,4L}(1, 4) &= -\frac{1}{2} \left[2p_k(1, 1)H_L(-3) + p_k(0, 1)H_L(-8) + 2p_k(\sqrt{2}, 1)H_L(-4) + p_k(0, 1)H_L(-16) \right. \\
&\quad \left. + 2p_k(2, 1)H_L(0) + 2B(L) - 6\mathbb{1}_{\substack{k=2 \\ L=\square}} \right].
\end{aligned}$$

Then grouping all the terms of $[*]$ by the $p_k(t, m)$ yields

$$\begin{aligned}
[*] &= \frac{1}{2} p_k(0, 1) \left[3H_L(-4) + H_{2L}(-4) - H_{4L}(-4) - H_L(-16) \right] \\
&\quad + p_k(1, 1) \left[2H_L(-3) + H_{2L}(-3) - H_{4L}(-3) \right] \\
&\quad + p_k(2, 1) \left[2H_L(0) + H_{2L}(0) - H_{4L}(0) \right] \\
&= \frac{1}{2} p_k(0, 1) \left[\frac{3}{2} \left(\frac{-4}{L} \right) - \frac{3}{2} \left(\frac{-16}{L} \right) \right] \\
&\quad + p_k(1, 1) \frac{\mathbb{1}_{3 \nmid L}}{3} \left[2 \left(\frac{-3}{L} \right) + \left(\frac{-3}{2L} \right) - \left(\frac{-3}{4L} \right) \right] \\
&\quad + p_k(2, 1) \frac{-1}{12} \left[2L + 2L - 4L \right]
\end{aligned}$$

$$= 0,$$

as desired. \square

5. μ_p -EQUIDISTRIBUTION OF HECKE EIGENVALUES

In this section, we prove Theorem 1.3. For a fixed prime p , let μ_p denote the p -adic Plancherel measure $\mu_p(x) = \frac{p+1}{\pi} \frac{(1-x^2/4)^{1/2}}{(p^{1/2} + p^{-1/2})^2 - x^2} dx$. Then over the spaces $S_k^\sigma(N)$ and $S_k^{\text{new},\sigma}(N)$ (for admissible sign patterns σ), we show that the Hecke eigenvalues for T'_p are μ_p -equidistributed as $N+k \rightarrow \infty$.

Theorem 1.3. *Fix p prime. Consider $N \geq 1$ coprime to p , σ sign patterns for N , and $k \geq 2$ even. Then the Hecke eigenvalues of $T'_p|_{S_k^\sigma(N)}$ are μ_p -equidistributed as $N+k \rightarrow \infty$. In other words, for any sequence $\{(N_i, k_i, \sigma_i)\}_{i \geq 1}$ with $N_i + k_i \rightarrow \infty$, the collection of Hecke eigenvalues $\text{eigv}(T'_m|_{S_{k_i}^{\sigma_i}(N_i)})$ is μ_p -equidistributed as $i \rightarrow \infty$. Moreover, if we assume it is not the case that $4 \parallel N$ with $\sigma(4) = +1$, then the Hecke eigenvalues of $T'_p|_{S_k^{\text{new},\sigma}(N)}$ are also μ_p -equidistributed as $N+k \rightarrow \infty$.*

Proof. We just state the argument here for $S_k^{\text{new},\sigma}(N)$, and an identical argument works for the full space $S_k^\sigma(N)$.

To show asymptotic μ_p -equidistribution of the Hecke eigenvalues, it suffices (e.g. by Kuipers-Niederreiter [10, Theorem 1.1], Serre [19, Proposition 2]) to show that

$$\frac{1}{\dim S_k^{\text{new},\sigma}(N)} \sum_{\lambda \in \text{eigv}\left(T'_p|_{S_k^{\text{new},\sigma}(N)}\right)} X_n(\lambda) \longrightarrow \int_{-2}^2 X_n(x) \mu_p(x) \quad \text{as } N+k \rightarrow \infty \quad (5.1)$$

for the Chebyshev polynomials $X_n(x) := U_n(x/2)$ for $n \geq 0$, since they span the space of all polynomials, which is a dense subspace of all continuous functions on $[-2, 2]$.

The integral on the right hand side of (5.1) has value $p^{-n/2} \mathbb{1}_{n \text{ even}}$, by Serre [19, Equation (20)].

Additionally, it is well-known (see [19, Lemme 1], for example) that

$$T'_{p^n} = X_n(T'_p).$$

This means that

$$\begin{aligned} \text{LHS of (5.1)} &= \frac{1}{\dim S_k^{\text{new},\sigma}(N)} \sum_{\lambda \in \text{eigv}\left(T'_p|_{S_k^{\text{new},\sigma}(N)}\right)} X_n(\lambda) \\ &= \frac{\text{Tr}_{S_k^{\text{new},\sigma}(N)} X_n(T'_p)}{\text{Tr}_{S_k^{\text{new},\sigma}(N)} T'_1} \\ &= \frac{\text{Tr}_{S_k^{\text{new},\sigma}(N)} T'_{p^n}}{\text{Tr}_{S_k^{\text{new},\sigma}(N)} T'_1} \\ &\rightarrow p^{-n/2} \mathbb{1}_{n \text{ even}} \quad \text{as } N+k \rightarrow \infty, \end{aligned}$$

where for the last step, we used Corollary 3.2 at $m = p^n$ and $m = 1$ (since σ is an admissible sign pattern). This verifies (5.1), completing the proof. \square

6. APPLICATIONS

Here, we state some applications of Theorem 1.3 analogous to those in Serre [19, §6].

For each $N \geq 1$ and σ admissible sign patterns for N , choose an eigenbasis $\{f_i\}_{i=1}^{\dim S_k^\sigma(N)}$ of $S_k^\sigma(N)$ with respect to the Hecke operators T_m satisfying $(m, N) = 1$. For each m coprime to N , let $x_{i,m}$ denote the T_m eigenvalue of f_i . Then for each i , let $K_i^\sigma := \mathbb{Q}(\{x_{i,m}\}_{(m,N)=1})$ denote the totally real field generated by the $x_{i,m}$ (see Cohen-Strömberg [6, Corollary 10.3.7(c), Remark 10.6.3(b)]). Then let $s(N, k, \sigma)_r$ be the number of i such that $[K_i^\sigma : \mathbb{Q}] = r$ (which is well-defined as the K_i^σ are independent of the eigenbasis chosen, up to permutation (cf. Serre, §6.1 [19])).

Similarly for $S_k^{\text{new}, \sigma}(N)$, we have the newform basis $\{f_i^{\text{new}}\}_{i=1}^{\dim S_k^{\text{new}, \sigma}(N)}$, and we define $K_i^{\text{new}, \sigma}$ and $s(N, k, \sigma)_r^{\text{new}}$ analogously. Then the following improves a result of Serre [19, Théorème 5] (and Binder [3, Corollary 10.0.5]), by extending to the sign pattern spaces.

Corollary 6.1. *Fix $k \geq 2$ even, p prime, and $r \geq 1$. Then for N coprime to p and σ admissible sign patterns for N ,*

$$\begin{aligned} \frac{s(N, k, \sigma)_r}{\dim S_k^\sigma(N)} &\rightarrow 0 & \text{as } N \rightarrow \infty, \\ \frac{s(N, k, \sigma)_r^{\text{new}}}{\dim S_k^{\text{new}, \sigma}(N)} &\rightarrow 0 & \text{as } N \rightarrow \infty. \end{aligned}$$

Proof. We show the proof only over the full space, as the proof over the newspace is identical. For N coprime to p , let $s(N, k, \sigma, p)_r$ denote the number of indices i such that $[\mathbb{Q}(x_{i,p}) : \mathbb{Q}] \leq r$. Note that $s(N, k, \sigma, p)_r \geq s(N, k, \sigma)_r$, so it suffices to show

$$\frac{s(N, k, \sigma, p)_r}{\dim S_k^\sigma(N)} \rightarrow 0$$

as $N \rightarrow \infty$ while $(N, p) = 1$. Note that if $x = x_{i,p}$ satisfies $[\mathbb{Q}(x_{i,p}) : \mathbb{Q}] \leq r$, then x also satisfies:

- (1) x is a totally real algebraic integer of degree $\leq r$;
- (2) For $\tau \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, the Galois conjugate x^τ of x satisfies $|x^\tau| \leq 2p^{(k-1)/2}$.

Let $A(p, k, r)$ be the set of all x satisfying conditions (1) and (2). Such a set must be finite, as the characteristic polynomials of its elements x must have degree $\leq r$ and bounded integer coefficients. Let $A' \subseteq [-2, 2]$ be the dilation of $A(p, k, r)$ under the mapping $x \mapsto \frac{x}{p^{(k-1)/2}}$. The set A' is finite, so it has μ_p measure 0, and it is independent of N and σ . Then since the $\frac{x_{i,p}}{p^{(k-1)/2}}$ are μ_p -equidistributed as $N \rightarrow \infty$ by Theorem 1.3, the proportion of i such that $\frac{x_{i,p}}{p^{(k-1)/2}} \in A'$ (and hence $x_{i,p} \in A$) tends to 0 as $N \rightarrow \infty$ (e.g. by Serre [19, Proposition 1]). This is equivalent to saying that $\frac{s(N, k, \sigma, p)_r}{\dim S_k^\sigma(N)} \rightarrow 0$ as $N \rightarrow \infty$, completing the proof. \square

We note here that over the sign pattern newspaces $S_k^{\text{new},\sigma}(N)$, we have $K_i^\sigma = \mathbb{Q}(f_i)$, where $\{f_i\}_{i=1}^{\dim S_k^{\text{new},\sigma}(N)}$ is the newform basis for $\dim S_k^{\text{new},\sigma}(N)$. From this perspective, Corollary 6.1 can be interpreted as a weak form of the Maeda Conjecture for general level.

Recall that the classical Maeda Conjecture states that $\deg \mathbb{Q}(f) = \dim S_k(1)$ (i.e. is maximum possible) for each normalized Hecke eigenform $f \in S_k(1)$. For squarefree levels N , a corresponding conjecture has also been stated for the newspaces. Since the Hecke operators restrict to the $S_k^{\text{new},\sigma}(N)$; for newforms $f \in S_k^{\text{new}}(N)$, the maximum possible degree of $\mathbb{Q}(f)$ is $\dim S_k^{\text{new},\sigma_f}(N)$, where σ_f denotes the sign pattern for f . In particular, Tsaknias and Chow-Ghitza posited a version of the Maeda Conjecture for squarefree level: that for newforms $f \in S_k^{\text{new}}(N)$, $\deg \mathbb{Q}(f) = \dim S_k^{\text{new},\sigma_f}(N)$ for sufficiently large weight k (see [21, Conjecture 2.4] and [4, Conjecture 5.2]).

For general level N , however, the picture is not as clear. In this case, there seem to be plenty of examples where $\deg \mathbb{Q}(f)$ is not maximal. However, weaker forms of the Maeda Conjecture have still been investigated. For example, Roberts conjectured that for any fixed weight $k \geq 6$, $\deg \mathbb{Q}(f) = 1$ for only finitely many newforms f [16, Conjecture 1.1]. Later, Martin made the same conjecture for arbitrary fixed degree r [12, Question 3]. Here, Corollary 6.1 can be interpreted as a density 1 version of these conjectures. In particular, for any fixed $r \geq 1$, Corollary 6.1 shows that the proportion of newforms $f \in S_k^{\text{new},\sigma}(N)$ with $\deg \mathbb{Q}(f) \leq r$ tends to 0 as $N \rightarrow \infty$ (along N coprime to p).

We can also use Corollary 6.1 to obtain information about the *largest* degree of K_i^σ and $K_i^{\text{new},\sigma}$. The following result follows immediately from Corollary 6.1.

Corollary 6.2 (cf. Serre, Théorème 6 [19]). *Fix $k \geq 2$ even and p prime. Let $r(N, k, \sigma)$ be the maximum degree among the K_i^σ associated with (N, k, σ) , where σ is an admissible sign pattern for N , and similarly define $r(N, k, \sigma)^{\text{new}}$. Then for N coprime to p and σ admissible sign patterns for N ,*

$$\begin{aligned} r(N, k, \sigma) &\rightarrow \infty & \text{as } N \rightarrow \infty, \\ r(N, k, \sigma)^{\text{new}} &\rightarrow \infty & \text{as } N \rightarrow \infty. \end{aligned}$$

Now, let $J_0(N)$ denote the Jacobian of the modular curve $X_0(N)$. Note that the Atkin-Lehner involutions act naturally on $X_0(N)$, and thus also act on $J_0(N)$ by functoriality. Up to \mathbb{Q} -isogeny, we have the decomposition:

$$J_0(N) \cong \prod_{\sigma} J_0^\sigma(N),$$

where $W_Q = \sigma(Q)$ on $J_0^\sigma(N)$ for all $Q \parallel N$. Now, each $J_0^\sigma(N)$ can be decomposed into a product of \mathbb{Q} -simple factors:

$$J_0^\sigma(N) \cong \prod A_j^\sigma.$$

Following the same argument as Serre (and Ribet [15]), we see that the number of factors A_j^σ with dimension r is exactly $\frac{1}{r}s(N, 2, \sigma)_r$ for each $r \geq 1$. Now, Corollary 6.2 implies the following.

Corollary 6.3 (cf. Serre, Théorème 7 [19]). *Fix p prime. Let $r(N, \sigma)$ be the largest dimension of the simple \mathbb{Q} -factors of $J_0^\sigma(N)$, where σ is an admissible sign pattern for N . Then for N coprime to p and σ admissible sign patterns for N ,*

$$r(N, \sigma) \rightarrow \infty \quad \text{as } N \rightarrow \infty.$$

ACKNOWLEDGEMENTS

We would like to thank Kimball Martin for his helpful comments on this paper. We would also like to thank the referee for many insightful comments, and especially for pointing out the reference [2] to Theorem 1.2.

This research was supported by NSF grant DMS-2349174. Research of Hui Xue is supported by Simons Foundation Grant MPS-TSM-00007911.

REFERENCES

- [1] Eran Assaf. A note on the trace formula. <https://arxiv.org/abs/2311.03523>, 2024.
- [2] A. O. L. Atkin and J. Lehner. Hecke operators on $\Gamma_0(m)$. *Math. Ann.*, 185:134–160, 1970.
- [3] John Binder. Fields of rationality of cusp forms. *Israel J. Math.*, 222(2):973–1028, 2017.
- [4] Sam Chow and Alexandru Ghitza. Distinguishing newforms. *Int. J. Number Theory*, 11(3):893–908, 2015.
- [5] Archer Clayton, Helen Dai, Tianyu Ni, Hui Xue, and Jake Zummo. Nonvanishing of second coefficients of hecke polynomials. *Journal of Number Theory*, 262:186–221, 2024.
- [6] Henri Cohen and Fredrik Strömberg. *Modular forms*, volume 179 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2017.
- [7] J. B. Conrey, W. Duke, and D. W. Farmer. The distribution of the eigenvalues of Hecke operators. *Acta Arith.*, 78(4):405–409, 1997.
- [8] Michael Griffin, Ken Ono, and Wei-Lun Tsai. Heights of points on elliptic curves over \mathbb{Q} . *Proc. Amer. Math. Soc.*, 149(12):5093–5100, 2021.
- [9] Henryk Iwaniec, Wenzhi Luo, and Peter Sarnak. Low lying zeros of families of L -functions. *Inst. Hautes Études Sci. Publ. Math.*, pages 55–131, 2000.
- [10] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1974.
- [11] Kimball Martin. Refined dimensions of cusp forms, and equidistribution and bias of signs. *J. Number Theory*, 188:1–17, 2018.
- [12] Kimball Martin. An on-average Maeda-type conjecture in the level aspect. *Proc. Amer. Math. Soc.*, 149(4):1373–1386, 2021.
- [13] Kimball Martin. Root number bias for newforms. *Proc. Amer. Math. Soc.*, 151(9):3721–3736, 2023.
- [14] Kimball Martin. Distribution of local signs of modular forms and murmurations of Fourier coefficients. *Mathematika*, 71(3), 2025.
- [15] Kenneth A. Ribet. Twists of modular forms and endomorphisms of abelian varieties. *Math. Ann.*, 253(1):43–62, 1980.
- [16] David P. Roberts. Newforms with rational coefficients. *The Ramanujan Journal*, 46(3):835–862, Aug 2018.
- [17] Erick Ross. Hecke eigenvalue equidistribution over the newspaces with nebentypus. *Submitted*, 2025.

- [18] Erick Ross. Proportion of Atkin-Lehner signs, 2025. <https://github.com/erick-ross/proportion-AL-signs>, 2025.
- [19] Jean-Pierre Serre. Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p . *J. Amer. Math. Soc.*, 10(1):75–102, 1997.
- [20] Nils-Peter Skoruppa and Don Zagier. Jacobi forms and a certain space of modular forms. *Invent. Math.*, 94(1):113–146, 1988.
- [21] Panagiotis Tsaknias. A possible generalization of Maeda's conjecture. In *Computations with Modular Forms*, pages 317–329. Springer International Publishing, 2014.

(E. Ross) SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, CLEMSON UNIVERSITY, CLEMSON, SC 29634
Email address: `erickjohnross@gmail.com`

(A. van Lidth) DEPARTMENT OF MATHEMATICS AND STATISTICS, AMHERST COLLEGE, AMHERST, MA 01002
Email address: `alexandrevanlidth@gmail.com`

(M. R. Wolf) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109
Email address: `martharose.wolf@gmail.com`

(H. Xue) SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, CLEMSON UNIVERSITY, CLEMSON, SC 29634
Email address: `huixue@clemson.edu`